

HOMOMORPHISM COMPLEXES AND K-CORES

GREG MALEN

ABSTRACT. We prove that the topological connectivity of a graph homomorphism complex $\text{Hom}(G, K_m)$ is at least $m - D(G) - 2$, where $D(G) = \max_{H \subseteq G} \delta(H)$. This is a strong generalization of a theorem of Čukić and Kozlov, in which $D(G)$ is replaced by the maximum degree $\Delta(G)$. It also generalizes the graph theoretic bound for chromatic number, $\chi(G) \leq D(G) + 1$, as $\chi(G) = \min\{m : \text{Hom}(G, K_m) \neq \emptyset\}$. Furthermore, we use this result to examine homological phase transitions in the random polyhedral complexes $\text{Hom}(G(n, p), K_m)$ when $p = c/n$ for a fixed constant $c > 0$.

1. INTRODUCTION

The study of graph homomorphism complexes, $\text{Hom}(G, H)$ for fixed graphs G and H , grew out of a rising interest in finding topological obstructions to graph colorings, as in Lovász's proof of the Kneser Conjecture [11]. Specifically, Lovász used the topological connectivity of the Neighborhood Complex, $\mathcal{N}(G)$, to provide a lower bound on the chromatic number of a Kneser graph. It turns out that $\mathcal{N}(G)$ is a specialization of a homomorphism complex, as it is homotopy equivalent to $\text{Hom}(K_2, G)$ (see Proposition 4.2 in [2]). The structure of the underlying graphs has many quantifiable effects on the topology of these complexes. The 0-cells of $\text{Hom}(G, H)$ are precisely the graph homomorphisms from $G \rightarrow H$, and so the most straightforward such result is that the chromatic number of G , $\chi(G)$, is the minimum m for which $\text{Hom}(G, K_m)$ is non-empty. For $\Delta(G)$ the maximum degree of G , Babson and Kozlov conjectured [2], and Čukić and Kozlov later proved [6] that if $\Delta(G) = d$, then $\text{Hom}(G, K_m)$ is at least $(m - d - 2)$ -connected. Note that when $m = d + 1$ this recovers the well known bound $\chi(G) \leq \Delta(G) + 1$.

In graph theory, there is an improvement of this bound using the minimum degree of induced subgraphs, namely that $\chi(G) \leq \max_{H \subseteq G} \delta(H) + 1$, for $\delta(H)$ the minimum degree of H . The value $D(G) := \max_{H \subseteq G} \delta(H)$ is known as the degeneracy of G , a graph property which is commonly utilized in computer science and the study of large networks. Using $D(G)$ in place of $\Delta(G)$, in Section 3 we prove a generalization of this graph theoretic result which will also specialize to the Čukić–Kozlov Theorem. In Section 4 we focus on the case that $H = K_3$ and show, with only a few notable exceptions, that $\text{Hom}(G, K_3)$ is disconnected if it is non-empty.

Furthermore, in the setting of random graphs the evolution of $D(G)$ for a sparse Erdős–Rényi graph $G \sim G(n, p)$ has been well studied. For $p = c/n$ for a constant $c > 0$, Pittel, Spencer, and Wormwald exhibited sharp thresholds for the appearance and size of subgraphs with minimum degree k [13]. In Section 5 we combine our results with their work to exhibit both phase transitions and lower bounds for the topological connectivity of the random polyhedral complexes $\text{Hom}(G(n, c/n), K_m)$ for constant $c > 0$. When $m = 3$,

we are able to use the results from Section 4 to evaluate the limiting probability for the connectivity of $\text{Hom}(G(n, c/n), K_3)$ for all $c > 0$.

2. BACKGROUND AND DEFINITIONS

In the following, all graphs are undirected simple graphs, and K_m is the complete graph on m vertices. For a graph G and a vertex $v \in V(G)$, when it is clear from context we may write simply that $v \in G$. In this section we define a graph homomorphism complex, or hom-complex, and various properties which will be used to acquire bounds on the topological connectivity of these complexes.

Definition 2.1. For fixed graphs G and H , a graph homomorphism from $G \rightarrow H$ is an edge preserving map on the vertices, $f : V(G) \rightarrow V(H)$ such that $\{f(v), f(u)\} \in E(H)$ if $\{v, u\} \in E(G)$.

Definition 2.2. For fixed graphs G and H , cells in $\text{Hom}(G, H)$ are functions

$$\eta : V(G) \rightarrow 2^{V(H)} \setminus \emptyset$$

with the restriction that if $\{x, y\} \in E(G)$, then $\eta(x) \times \eta(y) \subseteq E(H)$. The dimension of η is defined to be $\dim(\eta) = \sum_{v \in V(G)} (|\eta(v)| - 1)$, with ordering $\eta \subseteq \tau$ if $\eta(v) \subseteq \tau(v)$ for all $v \in V(G)$.

The set of 0-cells of $\text{Hom}(G, H)$ is then precisely the set of graph homomorphisms from $G \rightarrow H$, with higher dimensional cells formed over them as multihomomorphisms. Furthermore, every cell in a hom-complex is a product of simplices, and hom-complexes are entirely determined by their 1-skeletons. If η is a product of simplices and its 1-skeleton is contained in $\text{Hom}(G, H)$, then $\eta \in \text{Hom}(G, H)$. The 1-skeleton of $\text{Hom}(G, H)$ can be thought of in the following manner. Two graph homomorphisms $\eta, \tau : G \rightarrow H$ are adjacent in $\text{Hom}(G, H)$ if and only if their images differ on exactly one vertex $v \in G$. So $\eta(v) \neq \tau(v)$, $\eta(u) = \tau(u)$ for all $u \in G \setminus \{v\}$, and the 1-cell joining η and τ is the multihomomorphism σ for which $\sigma(v) = \eta(v) \cup \tau(v)$ and $\sigma(u) = \eta(u) = \tau(u)$ for all $u \in G \setminus \{v\}$.

Since the property of being a homomorphism is independent on disjoint connected components of a graph, hom-complexes obey a product rule for disjoint unions. For any graphs G, G' and H ,

$$\text{Hom}(G \amalg G', H) = \text{Hom}(G, H) \times \text{Hom}(G', H)$$

Thus, when convenient we can always restrict our attention to connected graphs. Here we also introduce the notion of a graph folding.

Definition 2.3. Denote the neighborhood of a vertex $v \in G$ by $N(v) = \{w \in G : w \sim v\}$. Let $v, u \in G$ be distinct vertices such that $N(v) \subseteq N(u)$. Then a fold of G is a homomorphism from $G \rightarrow G \setminus \{v\}$ which sends $v \mapsto u$ and is otherwise the identity.

Lemma 2.4. (Babson, Kozlov; Proposition 5.1 in [2]). If G and H are graphs and $v, u \in G$ are distinct vertices such that $N(v) \subseteq N(u)$, then the folding $G \rightarrow G \setminus \{v\}$ which sends $v \mapsto u$ induces a homotopy equivalence $\text{Hom}(G \setminus \{v\}, H) \rightarrow \text{Hom}(G, H)$.

If T is a tree, for example, then T folds to a single edge, so $\text{Hom}(T, K_m) \simeq \text{Hom}(K_2, K_m)$. And by Proposition 4.5 in [2], $\text{Hom}(K_n, K_m)$ is homotopy equivalent to a wedge of $(m - n)$ -spheres, and in particular $\text{Hom}(K_2, K_m) \simeq S^{m-2}$. So $\text{Hom}(T, K_m) \simeq S^{m-2}$ for any tree T . For a thorough introduction to hom-complexes, see [2].

Notation 2.5. For a graph G define the maximum degree to be $\Delta(G) := \max_{v \in V(G)} \{\deg(v)\}$, and the minimum degree to be $\delta(G) := \min_{v \in V(G)} \{\deg(v)\}$.

Definition 2.6. The k -core of a graph G is the subgraph $c_k(G) \subseteq G$ obtained by the process of deleting vertices with degree less than k , along with all incident edges, one at a time until there are no vertices with degree less than k .

Regardless of the order in which vertices are deleted, this process always terminates in the unique induced subgraph $c_k(G) \subseteq G$ which is maximal over all subgraphs of G which have minimum degree at least k . The k -core of a graph may be the empty graph, and the existence of a non-empty k -core is a monotone question, as $c_l(G) \subseteq c_j(G)$ whenever $j \leq l$. Here we are interested in the most highly connected non-trivial subgraph.

Definition 2.7. The degeneracy of G is $D(G) := \max_{H \subseteq G} \delta(H)$, for H an induced subgraph.

Then $D(G)$ is the maximum k such that $c_k(G)$ is non-empty. Given these definitions, we are now able to state the main theorem:

Theorem 2.8. For any graph G , $\text{Hom}(G, K_m)$ is at least $(m - D(G) - 2)$ -connected.

And $D(G) \leq \Delta(G)$, hence this will imply the Čukić–Kozlov Theorem. Our interest is primarily in applying this result to the case that G is a sparse Erdős–Rényi random graph, where $D(G)$ has been studied extensively and is much smaller than $\Delta(G)$. It should also be noted that Engström gave a similar strengthening of the Čukić–Kozlov Theorem in [8], replacing $\Delta(G)$ with a graph property which is independent of $D(G)$ and which may provide a better tool for studying the dense regime.

3. PROOF OF THEOREM 2.8

To prove the theorem, we first introduce a result of Csorba which finds subcomplexes that are homotopy equivalent to $\text{Hom}(G, K_m)$ by removing independent sets of G .

Notation 3.1. Let G and H be graphs. Define

$$\text{Ind}(G) := \{S \subseteq V(G) : S \text{ is an independent set}\}$$

And for $I \in \text{Ind}(G)$ define

$$\text{Hom}_I(G, H) := \{\eta \in \text{Hom}(G \setminus I, H) : \text{there is } \bar{\eta} \in \text{Hom}(G, H) \text{ with } \bar{\eta}|_{G \setminus I} = \eta\}$$

So $\text{Hom}_I(G, H)$ is the subcomplex of $\text{Hom}(G \setminus I, H)$ comprised of all multihomomorphisms from $G \setminus I \rightarrow H$ which extend to multihomomorphisms from $G \rightarrow H$.

Theorem 3.2. (Csorba; Theorem 2.36 in [5]) For any graphs G and H , and any $I \in \text{Ind}(G)$, $\text{Hom}(G, H)$ is homotopy equivalent to $\text{Hom}_I(G, H)$.

The proof of this is an application of both the Nerve Lemma and the Quillen Fiber Lemma, and a more thorough examination of this property is given by Schultz in [14]. Note that when $H = K_m$ and $I = \{v\}$ for any vertex $v \in G$, a multihomomorphism $\eta : G \setminus \{v\} \rightarrow K_m$

has an extension $\bar{\eta} : G \rightarrow K_m$ as long as there is some vertex in K_m which is not in $\eta(w)$ for any $w \in N(v)$. Thus

$$\text{Hom}_{\{v\}}(G, K_m) = \left\{ \eta \in \text{Hom}(G \setminus \{v\}, K_m) : \left| \bigcup_{w \in N(v)} \eta(w) \right| \leq m - 1 \right\}$$

Lemma 3.3. *Let G be a graph with a vertex v such that $\deg(v) \leq k$. Then the $(m - k - 1)$ -skeleton of $\text{Hom}(G \setminus \{v\}, K_m)$ is contained in $\text{Hom}_{\{v\}}(G, K_m)$.*

Proof of Lemma 3.3. Let $\eta \in \text{Hom}(G \setminus \{v\}, K_m) \setminus \text{Hom}_{\{v\}}(G, K_m)$. Then

$$m = \left| \bigcup_{w \in N(v)} \eta(w) \right| \leq \sum_{w \in G \setminus \{v\}} |\eta(w)|$$

And

$$\dim(\eta) = \sum_{w \in G \setminus \{v\}} (|\eta(w)| - 1) = \left(\sum_{w \in G \setminus \{v\}} |\eta(w)| \right) - |N(v)| \geq m - k$$

Hence if $\eta \in \text{Hom}(G \setminus \{v\}, K_m)$ with $\dim(\eta) \leq m - k - 1$, then $\eta \in \text{Hom}_{\{v\}}(G, K_m)$. $\square$

So if $\text{Hom}(G \setminus \{v\}, K_n)$ is $(m - k - 2)$ -connected, then $\text{Hom}_{\{v\}}(G, K_n)$ is $(m - k - 2)$ -connected. Combining this with Csorba's theorem, we have the following corollary and the proof of Theorem 2.8:

Corollary 3.4. *Let G be a graph with a vertex v such that $\deg(v) \leq k$. If $\text{Hom}(G \setminus \{v\}, K_n)$ is $(m - k - 2)$ -connected, then $\text{Hom}(G, K_n)$ is $(m - k - 2)$ -connected.*

Proof of Theorem 2.8. Let G be a graph with $D(G) = k$. Then $c_{k+1}(G)$ is the empty graph, and there is a sequence $G = G_0$, $G_i = G_{i-1} \setminus \{v_i\}$, with $\deg_{G_{i-1}}(v_i) \leq k$, terminating in $G_{|V(G)|} = \emptyset$. So $G_{|V(G)|-1}$ is a single vertex, and $\text{Hom}(G_{|V(G)|-1}, K_n) = \Delta^m$, the m -simplex, which is contractible, and thus $(m - k - 2)$ -connected. Hence, by induction, $\text{Hom}(G, K_m)$ is also $(m - k - 2)$ -connected. $\square$

4. $\text{Hom}(G, K_3)$

When G is a graph such that $\chi(G) \leq 3$, $\text{Hom}(G, K_3)$ has a particularly nice structure. If G does not have an isolated vertex, then $\text{Hom}(G, K_3)$ is a cubical complex, and Babson and Kozlov showed that it admits a metric with nonpositive curvature [2]. Notice that for a connected graph G with $\chi(G) \leq 3$, the bound obtained by Theorem 2.8 when $m = 3$ provides no new information. When $D(G) = 1$, G is a tree and folds to a single edge, so $\text{Hom}(G, K_3) \simeq \text{Hom}(K_2, K_3) \simeq S^1$. When $D(G) = 2$, the bound merely confirms that $\text{Hom}(G, K_3)$ is non-empty, and for $D(G) > 2$ it gives no information at all. Here we show that $\text{Hom}(G, K_3)$ is, in fact, disconnected for a large class of graphs G with $\chi(G) \leq 3$.

Theorem 4.1. *If G is a graph with $\chi(G) = 3$, then $\text{Hom}(G, K_3)$ is disconnected.*

This result has been formulated previously in the language of statistical physics, where Glauber dynamics examines precisely the 1-skeleton of $\text{Hom}(G(n, p), K_m)$. See, for example, the work of Cereceda, van den Heuvel and Johnson [3]. We provide a proof here in the context of work on hom-complexes of cycles done by Ćukić and Kozlov, and offer a

more general approach for lifting disconnected components via subgraphs in the following lemma.

Lemma 4.2. *Let $G \subseteq G'$ and H be graphs such that $\text{Hom}(G, H)$ is non-empty and disconnected. Let $\eta_1, \eta_2 \in \text{Hom}(G, H)$ be 0-cells, i.e. graph homomorphisms from $G \rightarrow H$, such that they are in distinct connected components of $\text{Hom}(G, H)$. If there are extensions $\overline{\eta}_1, \overline{\eta}_2 \in \text{Hom}(G', H)$ with $\overline{\eta}_i|_G = \eta_i$ for $i \in \{1, 2\}$, then $\text{Hom}(G', H)$ is also disconnected.*

Proof of Lemma 4.2. Let η_1, η_2 be graph homomorphisms from $G \rightarrow H$ such that they are in distinct connected components of $\text{Hom}(G, H)$, with extensions $\overline{\eta}_1, \overline{\eta}_2 \in \text{Hom}(G', H)$. We may assume that $\dim(\overline{\eta}_i) = 0$ for each i , since otherwise any 0-cells they contain would also be extensions of η_i . Suppose that $\overline{\eta}_1, \overline{\eta}_2$ are in the same connected component of $\text{Hom}(G', H)$. Then there is a path $\overline{\eta}_1 = \overline{\tau}_0 \sim \overline{\tau}_1 \sim \overline{\tau}_2 \sim \dots \sim \overline{\tau}_l = \overline{\eta}_2$ in $\text{Hom}(G', H)$, with $\dim(\overline{\tau}_j) = 0$ for all j . Let $\tau_j = \overline{\tau}_j|_G$, for $0 \leq j \leq l$. Then as functions on $V(G')$, for each j there is one vertex $v_j \in G'$ for which $\overline{\tau}_j(v_j) \neq \overline{\tau}_{j+1}(v_j)$, and they agree on all other vertices. If $v_j \in G' \setminus G$, then for the restrictions $\tau_j = \tau_{j+1}$. If $v_j \in G$, then $\tau_j(v_j) \neq \tau_{j+1}(v_j)$, but they agree on all other vertices, so $\tau_j \sim \tau_{j+1}$ in $\text{Hom}(G, H)$. Hence the path in $\text{Hom}(G', H)$ projects onto a possibly shorter path from η_1 to η_2 in $\text{Hom}(G, H)$, which is a contradiction.

Therefore the extensions $\overline{\eta}_1$ and $\overline{\eta}_2$ must be in different connected components of $\text{Hom}(G', H)$, which is thus disconnected. $\square$

The subgraphs we examine to obtain disconnected components of $\text{Hom}(G, K_3)$ will be cycles. In [7], Čukić and Kozlov fully characterized the homotopy type of $\text{Hom}(C_n, C_m)$ for all $n, m \in \mathbb{N}$. In particular, they showed that all 0-cells in a given connected component have the same number of what they call return points. For these complexes, let $V(C_n) = \{v_1, \dots, v_n\}$ such that $v_i \sim v_{(i+1) \bmod n}$ for all $1 \leq i \leq n$, and let $V(C_m) = \{1, \dots, m\}$ such that $j \sim (j+1) \bmod m$ for all $1 \leq j \leq m$. For the purpose of defining the return number of a 0-cell η , we momentarily drop the set bracket notation and write $\eta(v_i) = j$.

Definition 4.3. *A return point of a 0-cell $\eta \in \text{Hom}(C_n, C_m)$ is a vertex $v_i \in C_n$ such that $\eta(v_{i+1}) - \eta(v_i) \equiv -1 \bmod m$. Then $r(\eta)$, the return number of η , is the number of $v_i \in C_n$ which are return points of η .*

Note that since η is a homomorphism, the quantity $\eta(v_{i+1}) - \eta(v_i)$ is always $\pm 1 \bmod m$. It simply measures in which direction C_n is wrapping around C_m on the edge $\{v_i, v_{i+1}\}$.

Lemma 4.4. (Čukić, Kozlov, Lemma 5.3 in [7]) *If two 0-cells of $\text{Hom}(C_n, C_m)$ are in the same connected component, then they have the same return number.*

For a 0-cell $\eta \in \text{Hom}(G, K_m)$, one may always obtain another 0-cell by swapping the inverse images of two vertices in K_m . When the target graph is a complete graph, we refer to the inverse image of a vertex in K_m as a color class of η . When $m = 3$, we have $K_3 = C_3$ and we can track the effect that interchanging two color classes has on the return number.

Notation 4.5. *For a 0-cell $\eta \in \text{Hom}(G, K_m)$, let $\eta_{[l,j]}$ denote the (l, j) -interchanging 0-cell obtained by defining $\eta_{[l,j]}^{-1}(l) = \eta^{-1}(j)$, $\eta_{[l,j]}^{-1}(j) = \eta^{-1}(l)$, and $\eta_{[l,j]}^{-1}(i) = \eta^{-1}(i)$ for all $i \notin \{l, j\}$.*

Lemma 4.6. *Fix a pair $\{l, j\} \subset \{1, 2, 3\}$, $l \neq j$. Then for any 0-cell $\eta \in \text{Hom}(C_n, K_3)$, every $v_i \in C_n$ is a return point for exactly one of η and $\eta_{[l,j]}$. Hence $r(\eta_{[l,j]}) = n - r(\eta)$.*

Proof of Lemma 4.6. Consider the edge $\{v_i, v_{i+1}\}$. Since the target graph is K_3 , we have that $\{\eta(v_i), \eta(v_{i+1})\} \cap \{l, j\} \neq \emptyset$. If $\{\eta(v_i), \eta(v_{i+1})\} = \{l, j\}$, then

$$\eta_{\{l,j\}}(v_{i+1}) - \eta_{\{l,j\}}(v_i) = -(\eta(v_{i+1}) - \eta(v_i))$$

Thus v_i is a return point of η if and only if it is not a return point of $\eta_{\{l,j\}}$. Alternatively, if $|\{\eta(v_i), \eta(v_{i+1})\} \cap \{l, j\}| = 1$, then one of v_i and v_{i+1} has a stationary image under an (l, j) -interchange while the other does not. By fixing the image of one vertex and changing the other, this flips the direction that C_n is wrapping around K_3 on the edge $\{v_i, v_{i+1}\}$. So returns become non-returns and vice versa. Therefore each $v_i \in C_n$ is a return point of exactly one of η and $\eta_{\{l,j\}}$ for any fixed $\{l, j\} \subset \{1, 2, 3\}$. $\square$

The proof of Theorem 4.1 will then proceed by finding an odd cycle in G and using a color class interchange to produce 0-cells in distinct connected components, which can be lifted from $\text{Hom}(C_{2k+1}, K_3)$ to $\text{Hom}(G, K_3)$.

Proof of Theorem 4.1 Since $\chi(G) = 3$, $\text{Hom}(G, K_3)$ is non-empty and G contains an odd cycle $H = C_{2k+1}$ for some $k \in \mathbb{N}$. Note that we do not require H to be an induced cycle. Let $\eta \in \text{Hom}(G, K_3)$ be a 0-cell, and let $\tau = \eta|_H$ be the induced homomorphism on H . Fix a distinct pair $\{l, j\} \subset \{1, 2, 3\}$ and consider the (l, j) -interchange $\eta_{\{l,j\}}$. It is straightforward that $\eta_{\{l,j\}}|_H = \tau_{\{l,j\}} \in \text{Hom}(H, K_3)$. Thus by Lemma 4.6, $r(\tau_{\{l,j\}}) = 2k + 1 - r(\tau) \neq r(\tau)$. Hence, by lemma 4.4 they are in different connected components of $\text{Hom}(H, K_3)$, and by Lemma 4.2 their extensions η and $\eta_{\{l,j\}}$ are in different connected components of $\text{Hom}(G, K_3)$, which is thus disconnected. $\square$

Bipartite graphs are a bit more complicated. We have seen that for any graph G which folds to a single edge, $\text{Hom}(G, K_3) \simeq S^1$. When G is bipartite and does not fold to an edge, it must contain an even cycle C_{2k} for $k \geq 3$. So by the same method one can start with $\eta \in \text{Hom}(G, K_3)$ and take its restriction $\eta|_{C_{2k}} = \tau \in \text{Hom}(C_{2k}, K_3)$. But if $r(\tau) = k$, then interchanging two color classes no longer guarantees disjoint components. In particular, the existence of too many 4-cycles close to C_{2k} may force $r(\tau) = k$. For example, for Q_3 the 1-skeleton of the 3-cube, $\text{Hom}(Q_3, K_3)$ is connected. Note that $D(Q_3) = 3$, so the lower bound achieved by Theorem 2.8 does not provide any information. Restricting to the case that G omits the subgraphs in Figure 4.1 is sufficient to ensure that this does not happen, and that $\text{Hom}(G, K_3)$ is disconnected.




FIGURE 4.1. If $\text{Hom}(G, K_3)$ is connected, then G must contain H_1 or H_2 .

Theorem 4.7. *If G is a finite, bipartite, connected graph which does not fold to an edge and does not contain H_1 or H_2 as a subgraph, then $\text{Hom}(G, K_3)$ is disconnected.*

To show this requires a great deal of case-by-case structural analysis when a minimal even cycle in G has length 6, 8, or 10. The method of the proof will suggest a stronger, albeit more technical result.

Proof of Theorem 4.7. Let G be a finite, bipartite, connected graph which does not fold to an edge and does not contain H_1 or H_2 as a subgraph. Since G is finite, bipartite and does not fold to an edge, it must contain an even cycle of length greater than 4. Let $H = C_{2k}$, $k \geq 3$ be a fixed cycle in G which has minimal length over all cycles in G of length greater than 4. Label the vertices of H as $v_1, \dots, v_{2k}$ such that $v_i \sim v_{(i+1) \bmod 2k}$ for all i . H cannot have internal chords, as any such edge would create a cycle C_l for $4 < l < 2k$, or in the case that $k = 3$ an antipodal chord would create a copy of H_1 . For $2k \equiv i \bmod 3$, take $\tau \in \text{Hom}(H, K_3)$ such that $\tau(v_j) = \{j \bmod 3\}$ for $j \leq 2k - i$, with $\tau(v_{2k}) = \{2\}$ if $i = 1$, and $\tau(v_{2k-1}) = \{1\}$, $\tau(v_{2k}) = \{2\}$ if $i = 2$. Figure 4.2 depicts τ for $k \in \{3, 4, 5\}$.




FIGURE 4.2. Image of τ on $H = C_{2k}$

As defined, only the last two vertices of C_{2k} can be return points of τ . So $r(\tau) \leq 2$ in all cases, and hence is not equal to k for any $k \geq 3$. Then for any $\{l, j\} \subset \{1, 2, 3\}$, $r(\tau_{\{l, j\}}) \neq r(\tau)$. Thus $\tau_{\{l, j\}}$ and τ are in distinct connected components of $\text{Hom}(H, K_3)$. So if we can construct an extension η to all of G , then $\eta_{\{l, j\}}$ will extend $\tau_{\{l, j\}}$ and hence η and $\eta_{\{l, j\}}$ will be in distinct connected component of $\text{Hom}(G, K_3)$.

Define $d(u, v)$ to be the minimal number of edges in a path connecting two vertices $u, v \in G$. Let σ be a bipartition of G , viewed as a 0-cell of $\text{Hom}(G, K_3)$ with image contained in $\{1, 2\}$. We will let $\eta = \sigma$ on $G \setminus H$ and then make adjustments as necessary when the definition conflicts with τ . Define the following sets:

$$\begin{aligned}
 B_i &:= \{u \in G : \min\{d(u, v) : v \in H\} = i\} \\
 A &:= \{u \in B_1 : \exists v \in N(u) \cap H \text{ with } \sigma(u) = \tau(v)\} \\
 A_{i,j} &:= \{u \in A : |N(u) \cap H| = i, |N(u) \cap A| = j\}
 \end{aligned}$$

So B_i is the set of points distance i from H , and A is the set of points in $G \setminus H$ on which we cannot define $\eta = \sigma$. To arrive at an appropriate definition for η on the set A , we must first make some structural observations concerning the partitioning of A into the subsets $A_{i,j}$.

Claim 4.8. *For every $u \in B_1$, $|N(u) \cap H| \leq 2$. And if $|N(u) \cap H| = 2$, then $N(u) \cap H = \{v_i, v_{(i+2) \bmod 2k}\}$ for some $i \leq 2k$.*

Proof of Claim 4.8. Let $u \in B_1$. Then connecting u to any two points in H creates a cycle of length at most $k + 2 < 2k$, as in Figure 4.3 (a). Since G is bipartite and H is minimal with respect to having length at least 6, this is valid only if it creates a 4-cycle, with

$N(u) \cap H = \{v_i, v_{(i+2) \bmod 2k}\}$ for some $i \leq 2k$. Similarly, any further points of attachment would need to be in a 4-cycle with each of the other two vertices in $N(u) \cap H$. But this is only possible if $H = C_6$, in which case it would create several copies of H_1 , shown in Figure 4.3 (b). $\square$




FIGURE 4.3. Obstructions to $|N(u) \cap H| > 2$

This imposes that $i \leq 2$ for $A_{i,j} \subseteq A$, while the following two claims will show that j is at most 1, and that $j = 1$ only if $i = 1$.

Claim 4.9. *Let $k \geq 4$, and let $u \in B_1$. Then $|N(u) \cap B_1| \leq 1$, and if $N(u) \cap B_1 \neq \emptyset$ then $|N(u) \cap H| = 1$.*

Proof of Claim. Let $k \geq 4$, let $u \in B_1$, and let $w \in N(u) \cap B_1$, with $u \sim v_i$ and $w \sim v_j$. G is bipartite, so $v_i \neq v_j$. If $v_i \not\sim v_j$, as in Figure 4.4 (a), there is a cycle of length l with $4 < l \leq k + 3 < 2k$. So $v_i \sim v_j$. Suppose further that $|N(u) \cap H| = 2$. Then $N(u) \cap H = \{v_{(j-1) \bmod 2k}, v_{(j+1) \bmod 2k}\}$ since both must be adjacent to v_j , as in Figure 4.4 (b). But then the induced graph on $\{u, w, v_{(j-1) \bmod 2k}, v_j, v_{(j+1) \bmod 2k}\}$ is a copy of H_2 . So if $N(u) \cap B_1 \neq \emptyset$ for $u \in B_1$, then $|N(u) \cap H| = 1$. Now suppose $|N(u) \cap B_1| > 1$. Then there is a vertex $x \in N(u) \cap B_1$ with $x \neq w$. As with w , the vertex in $N(x) \cap H$ must be adjacent to v_i . If $x \sim v_j$, as in Figure 4.4 (c), then $\{u, w, x, v_i, v_j\}$ is a copy of H_2 , and if x is connected to the other neighbor of v_i , as in Figure 4.4 (d), then it creates a copy of H_1 . Thus $|N(u) \cap B_1| \leq 1$. $\square$




FIGURE 4.4. Invalid structures in G if Claim 4.9 is false.

For $k = 3$ a similar result holds when B_1 is replaced by the subset A :

Claim 4.10. *Let $k = 3$, and let $u \in A$. Then $|N(u) \cap A| \leq 1$, and if $N(u) \cap A \neq \emptyset$ then $|N(u) \cap H| = 1$.*

Proof of Claim. Let $u \in A$ and let $w \in N(u) \cap A$. Recall that for $2k = 6$, $\tau(v_i) = \{i \bmod 3\}$ for all i . Then without loss of generality assume that $\sigma(u) = \{1\}$ and $u \sim v_1$. So $\sigma(w) = \{2\}$, and since $w \in A$ it must be adjacent to at least one of v_2 and v_5 . But G is bipartite and $u \sim v_5$ creates a 5-cycle, as indicated in Figure 4.5 (a). Thus $w \sim v_2$, and the same argument holds to show that $u \sim v_1$ if one first assumes that $w \sim v_2$. Furthermore, if $w \sim v_4$ or $w \sim v_6$ there is a copy of H_1 or H_2 , respectively, depicted in Figure 4.5 (b), (c). By symmetry $u \not\sim v_3$, $u \not\sim v_5$. Hence, if $u \in A$ with $N(u) \cap A \neq \emptyset$, then $|N(u) \cap H| = 1$. Now suppose there is $x \in N(u) \cap A$ with $x \neq w$. Then $\sigma(x) = \{2\}$, and by the preceding argument we have that $x \sim v_2$. But, as shown in Figure 4.5 (d), $\{u, w, x, v_1, v_2\}$ is then a copy of H_2 . Hence for $u \in A$, $|N(u) \cap A| \leq 1$. $\square$




FIGURE 4.5. Invalid structures in G if Claim 4.10 is false.

Then for all $k \geq 3$ we have that $A = A_{1,0} \cup A_{2,0} \cup A_{1,1}$, and vertices in $A_{1,1}$ which are adjacent to each other have adjacent attaching vertices in H . When possible, we will define $\eta(v) = \{3\}$ for vertices $v \in A$. However, this is not possible on all vertices in $A_{1,1}$, nor for vertices in $A_{2,0}$ which have a neighbor in H that is already assigned $\{3\}$. So define

$$\bar{A} = \{u \in A_{2,0} : \{\tau(v) : v \in N(u) \cap H\} = \{\sigma(u), \{3\}\}\} \cup \{u \in A_{1,1} : \sigma(u) = \{2\}\}$$

Now define $\eta : G \rightarrow K_3$ by:

$$\eta(u) = \left\{ \begin{array}{ll} \tau(u) & \text{if } u \in H \\ \{1\} & \text{if } u \in \bar{A} \cap A_{1,1}; \\ & u \in \bar{A} \cap A_{2,0} \text{ with } \sigma(u) = \{2\} \\ \{2\} & \text{if } u \in \bar{A} \cap A_{2,0} \text{ with } \sigma(u) = \{1\} \\ \{3\} & \text{if } u \in A \setminus \bar{A}; \\ & u \in G \setminus (A \cup H) \text{ with } N(u) \cap \bar{A} \neq \emptyset \\ \sigma(v) & \text{else} \end{array} \right\}$$

Notice that for $u \in \bar{A}$, $\eta(u) = \{1, 2\} \setminus \sigma(u) = \sigma(w)$ for any $w \in N(u)$. Hence the last qualifier assigns $\{3\}$ to any neighbors of vertices in $\bar{A}$ which need to be switched, and $\eta = \sigma$ on all remaining vertices.

It is clear that this defines a homomorphism on $A \cup H$, and on any edges between vertices $u \in G \setminus (A \cup H)$ for which $N(u) \cap \bar{A} = \emptyset$. Thus it suffices to check that this is consistent on edges with a vertex $u \in G \setminus (A \cup H)$ for which $N(u) \cap \bar{A} \neq \emptyset$. Note that by definition, any such u will have $\eta(u) = \{3\}$. So to ensure that $\eta(w) \neq \{3\}$ for any $w \in N(u)$,

we must check that $N(u) \cap (A \setminus \bar{A}) = \emptyset$, and for all $w \in N(u)$ that $N(w) \cap \bar{A} = \emptyset$. The cases $k = 3$ and $k \geq 4$ must again be handled separately.

Claim 4.11. *Let $k \geq 4$, and let $u \in G \setminus (A \cup H)$ with $x \in N(u) \cap \bar{A}$. Then $N(u) \cap (A \setminus \bar{A}) = \emptyset$, and if $w \in N(u)$ then $N(w) \cap \bar{A} = \emptyset$.*

Proof of Claim. Let u and x be as in the claim. Since $x \in A \subseteq B_1$, it must be that $u \in (B_1 \cup B_2) \setminus A$. Suppose first that $u \in B_1$. Then $|N(u) \cap B_1| \leq 1$ by Claim 4.9, and hence $N(u) \cap A = \{x\}$. So $N(u) \cap (A \setminus \bar{A}) = \emptyset$. Similarly, Claim 4.9 implies that $N(x) \cap B_1 = \{u\}$ and that $|N(x) \cap H| = 1$, so $x \in A_{1,0}$, which contradicts our assumption that $x \in \bar{A} \subseteq A_{2,0} \cup A_{1,1}$. So $u \notin B_1$.

Suppose instead that $u \in B_2$, and thus $N(u) \subseteq B_1 \cup B_2 \cup B_3$. For $w \in N(u) \cap B_3$ it is immediate that $w \notin A$ and that $(N(w) \cap \bar{A}) \subseteq (N(w) \cap B_1) = \emptyset$. Then assume that there is $w \in N(u) \cap B_1$, $w \neq x$. Then $\sigma(w) = \sigma(x)$ since G is bipartite, and thus for any attaching points in H , $v_i \sim x$, $v_j \sim w$, it must be that i and j are either both odd or both even. Since $x \in A_{2,0} \cup A_{1,1}$ we may assume that either $i \neq j$ or $x \in A_{1,1}$. Recall that an adjacent pair in $A_{1,1}$ forms a square with its adjacent pair in H . So if $i = j$ and $x \in A_{1,1}$ with $\{z\} = N(x) \cap A_{1,1}$, then z is adjacent to one of the two vertices in $N(v_i) \cap H$. As shown in Figure 4.6 (a), this creates a copy of H_1 . So we may assume that $i \neq j$. But then the short path between v_i and v_j in H along with $\{u, w, x\}$ creates a cycle of length l , $6 \leq l \leq k + 4$.




FIGURE 4.6. $w \in N(u) \cap B_1$, $w \neq x$.

If $k \geq 5$ this length is less than $2k$, as in Figure 4.6 (b), which contradicts the minimality of H . If $k = 4$, this creates a cycle of length 8 only if $j = (i + 4) \bmod 8$, as in Figure 4.6 (c). Recall that as defined for $k = 4$ (see Figure 4.2 (b)), $\tau(v_i) \neq \tau(v_{(i+4) \bmod 8})$ for all i . But $\sigma(x) = \sigma(w)$, and therefore x and w cannot both be in A . As this is true for any pair of vertices in $N(u) \cap B_1$, it must be that $|N(u) \cap A| \leq 1$, and hence by assumption $N(u) \cap A = \{x\}$. Furthermore, for any $w \in B_1 \setminus A$, if $w \sim z$ with $z \in A$, then by Claim 4.9 it must be that $z \in A_{1,0}$. Thus $N(w) \cap \bar{A} = \emptyset$ for all $w \in N(u) \cap B_1$.

Now suppose that $w \in N(u) \cap B_2$. Note that $w \notin A \subseteq B_1$. Then let $x = x_u$ and let $x_w \in N(w) \cap B_1$. Note that $\sigma(x_u) \neq \sigma(x_w)$ since there is a path of length 3 from x_u to x_w , so the connecting vertices for x_u and x_w in H must be distinct to avoid creating an odd cycle. Then the path between their connecting vertices in H along with x_u, u, w , and x_w creates a cycle of length l , $6 \leq l \leq k + 5$. For $k > 5$ such a cycle contradicts the minimality of H .

Suppose $k = 4$. Then to avoid creating a cycle of length less than 8, this construction requires that the path in H between the connecting points of x_u and x_w is length 3. Suppose by way of contradiction that $x_w \in \bar{A}$. So both $x_u, x_w \in \bar{A} \subseteq A_{2,0} \cup A_{1,1}$. Without loss of

generality, assume that $x_u \in A_{2,0}$ with $N(x_u) \cap H = \{v_i, v_{(i+2) \bmod 8}\}$ for some $1 \leq i \leq 8$. Then $v_{(i+5) \bmod 8}$ is the only vertex in H which is distance 3 from both v_i and $v_{(i+2) \bmod 8}$. So x_w cannot also be in $A_{2,0}$, and thus $x_w \in A_{1,1}$ with $N(x_w) \cap H = \{v_{(i+5) \bmod 8}\}$. By definition, $\sigma(v) = \{2\}$ for all $v \in \bar{A} \cap A_{1,1}$, so we have that $\sigma(x_w) = \{2\}$, and thus also $\tau(v_{(i+5) \bmod 8}) = \{2\}$. For τ as defined, this forces $i \in \{3, 5, 8\}$. But now $\sigma(x_u) = \{1\}$, and so $x_u \in \bar{A} \cap A_{2,0}$ implies that $\{\tau(v_i), \tau(v_{(i+2) \bmod 8})\} = \{\{1\}, \{3\}\}$, which is only possible if $i \in \{1, 4\}$. Figure 4.7 (a) and (b) shows the arrangements if $i = 1$ and $i = 4$ respectively, with $\tau(v_6), \tau(v_1) \neq \{2\}$. Hence it must be that both $x_u, x_w \in A_{1,1}$. However, $\sigma(x_u) \neq \sigma(x_w)$, and as noted above $\sigma(v) = \{2\}$ for all $v \in \bar{A} \cap A_{1,1}$. So at most one of x_u and x_w can be in $\bar{A}$, and by assumption $x_u \in \bar{A}$. Thus $x_w \notin \bar{A}$ and $N(w) \cap \bar{A} = \emptyset$ for all $w \in N(u) \cap B_2$.





FIGURE 4.7. $x_u \in \bar{A} \cap A_{2,0}, x_w \in \bar{A} \cap A_{1,1}, \sigma(x_u) = \{1\}$ when $k = 4$.

Now suppose $k = 5$, and again assume that $x_w \in \bar{A}$. Here cycles created by adjoining u, w, x_u , and x_w will be of length 10 only if the connecting vertices in $H = C_{10}$ are distance 5 apart. Hence neither x_u nor x_w can be in $A_{2,0}$, and both must be in $A_{1,1}$. Thus, as in the case above for $k = 4$, $\sigma(x_u) \neq \sigma(x_w)$ implies that at most one of x_u and x_w can be in $\bar{A}$. Therefore $x_w \notin \bar{A}$ and $N(w) \cap \bar{A} = \emptyset$ for all $w \in N(u) \cap B_2$. $\square$


Claim 4.12. *Let $k = 3$, and let $u \in G \setminus (A \cup H)$ with $x \in N(u) \cap \bar{A}$. Then $N(u) \cap (A \setminus \bar{A}) = \emptyset$, and if $w \in N(u)$ then $N(w) \cap \bar{A} = \emptyset$.*

Proof of Claim. Let vertices u and x be as in the claim, and let $v_i \in N(x) \cap H$ be a vertex such that $\sigma(x) = \tau(v_i)$, which exists by the assumption that $x \in A$. Then since G is bipartite, it must be that $\{v \in H : d(v, u) = 2\} \subseteq \{v_i, v_{(i+2) \bmod 6}, v_{(i+4) \bmod 6}\}$. Recall that τ is defined such that for each $j \in K_3$ the set $\tau^{-1}(\{j\})$ contains one vertex with an odd index and one vertex with an even index (see Figure 4.2 (a)). So $\{v_i, v_{(i+2) \bmod 6}, v_{(i+4) \bmod 6}\} \cap \tau^{-1}(\sigma(x)) = \{v_i\}$. Hence $w \sim v_i$ for every $w \in N(u) \cap A$. Then if $|N(u) \cap A| \geq 3$, this would create a copy of H_2 , depicted in Figure 4.8 (a). Thus $|N(u) \cap A| \leq 2$.


FIGURE 4.8. $k = 3$ and $|N(u) \cap A| \geq 2$.

Suppose that $|N(u) \cap A| = 2$. If there was an adjacency from the set $\{v_{(i+2) \bmod 6}, v_{(i+4) \bmod 6}\}$ to $N(u) \cap A$, it would create a copy of H_1 , as in Figure 4.8 (b). So $N(u) \cap A_{2,0} = \emptyset$. And adjacent pairs in $A_{1,1}$ form squares with their connecting pair in H . So if $w \in N(u) \cap A_{1,1}$, then there is $w' \in N(w) \cap A_{1,1}$ such that, without loss of generality, $w' \sim v_{(i+1) \bmod 6}$. But, as depicted in Figure 4.8 (c), this creates a copy of H_1 . Thus it cannot be that both $|N(u) \cap A| > 1$ and $N(u) \cap (A_{2,0} \cup A_{1,1}) \neq \emptyset$. Since $x \in \bar{A} \subseteq A_{2,0} \cup A_{1,1}$ by assumption, we have that $N(u) \cap A = \{x\}$, and so $N(u) \cap (A \setminus \bar{A}) = \emptyset$.


Now it suffices to check for all $w \in N(u) \setminus (H \cup \{x\})$ that $N(w) \cap \bar{A} = \emptyset$. Suppose that $u \in B_1$. Then $N(u) \cap H \subset \{v_{(i+1) \bmod 6}, v_{(i+3) \bmod 6}, v_{(i+5) \bmod 6}\}$ since $x \sim v_i$. Further suppose that $x \in A_{2,0}$, so without loss of generality $N(x) \cap H = \{v_i, v_{(i+2) \bmod 6}\}$. Then all three possible adjacencies for u in H are shown in Figure 4.9, with $u \sim v_{(i+1) \bmod 6}$ creating a copy of H_2 , while $v \sim v_{(i+3) \bmod 6}$ or $v \sim v_{(i+5) \bmod 6}$ each creates a copy of H_1 . Hence $x \notin A_{2,0}$.

FIGURE 4.9. $k = 3$, $u \in B_1$, and $x \in A_{2,0}$.

So $x \in \bar{A} \cap A_{1,1}$ which implies that $\sigma(x) = \{2\}$, and so $\sigma(u) = \{1\}$. Then without loss of generality let $i = 2$. Note $x \sim v_2$ implies u must be connected to a vertex in H which has odd index. But $u \notin A$, so $u \not\sim v_1$. Let $\{w\} = N(x) \cap A_{1,1}$, so it must be that $\sigma(w) = \{1\}$ and $w \sim v_1$. Then $u \sim v_3$ would create a copy of H_1 , shown in Figure 4.10 (a), and so $u \sim v_5$. Now consider a vertex $z \in N(u) \setminus (H \cup \{x\})$, and suppose there exists $v_z \in N(z) \cap \bar{A}$. Then $d(u, v_z) = 2$, so $\sigma(v_z) = \sigma(u) = \{1\}$. And by definition, we have that $\{v \in \bar{A} : \sigma(v) = \{1\}\} \subseteq A_{2,0}$, so $v_z \in A_{2,0}$ and the two vertices in $N(v_z) \cap H$ have images $\{1\}$ and $\{3\}$ under σ . Since $\sigma(v_z) = \sigma(u)$, vertices in $N(v_z) \cap H$ must have odd index. Thus $N(v_z) \cap H = \{v_1, v_3\}$. As shown in Figure 4.10 (b), $\{w, x, v_z, v_1, v_2, v_3\}$ is then a copy of H_1 . So if $u \in B_1$ there cannot be any vertex $z \in N(u)$ with $N(z) \cap \bar{A} \neq \emptyset$.

FIGURE 4.10. $k = 3$, $u \in B_1$, and $x \in \bar{A} \cap A_{1,1}$.

Now assume that $u \in B_2$. Let $x = x_u$, and suppose that there is a vertex $w \in N(u)$ which has a vertex $x_w \in N(w) \cap \bar{A}$. Then there is a path of length 3 from x_u to x_w , so $x_u \neq x_w$ and also $\sigma(x_u) \neq \sigma(x_w)$. Then they cannot both be in $A_{1,1}$, since $\sigma(v) = \{2\}$ for all $v \in \bar{A} \cap A_{1,1}$. Without loss of generality, let $x_u \in A_{2,0}$. Suppose that $x_w \in A_{1,1}$ and let $\{z\} = N(x_w) \cap A_{1,1}$. Then $\sigma(x_w) = \{2\}$ and without loss of generality we have that $x_w \sim v_2$ and $z \sim v_1$. Now x_u must be adjacent to two vertices in H with odd indices, one of which has image $\{3\}$ under τ . And $\sigma(x_u) = \{1\}$, so $N(x_u) \cap H = \{v_1, v_3\}$. This is depicted in Figure 4.11 (a), where $\{v_1, v_2, v_3, x_u, x_w, z\}$ forms a copy of H_1 . Suppose instead that $x_w \in A_{2,0}$. Then each x_w and x_u must be adjacent to a vertex whose image under τ is $\{3\}$. Without loss of generality assume that $x_u \sim v_3$ and $x_w \sim v_6$, and let $x_u \sim v_j$ and $x_w \sim v_l$ be the other two adjacencies. Then j must be odd and l must be even, and $\tau(v_j) = \sigma(x_u) \neq \sigma(x_w) = \tau(v_l)$. So either $j = 1$ and $l = 2$, or $j = 5$ and $l = 4$. By symmetry we may assume that $x_u \sim v_1$ and $x_w \sim v_2$, as in Figure 4.11 (b). But then $\{v_1, v_2, v_3, v_6, x_u, x_w\}$ forms a copy of H_1 . Thus $N(w) \cap \bar{A} = \emptyset$ for all $w \in N(u)$. $\square$

FIGURE 4.11. $k = 3$, $u \in B_2$, and $\{x_u, x_w\} \cap A_{2,0} \neq \emptyset$.

It follows immediately from Claims 4.11 and 4.12 that $\eta : G \rightarrow K_3$ is a homomorphism for all $k \geq 3$. And η extends τ , so $\eta_{[l,j]}$ extends $\tau_{[l,j]}$. Thus by Lemma 4.2, η and $\eta_{[l,j]}$ are in distinct connected components of $\text{Hom}(G, K_3)$, which is therefore disconnected. $\square$

The details of this proof suggests that it is only necessary to require that there be some cycle $C_{2k} \subset G$, $k \geq 3$, such that $H' \cap C_{2k} = \emptyset$ for any subgraph $H' \subset G$ with $H' \in \{H_1, H_2\}$. But even this leaves out many bipartite G for which $\text{Hom}(G, K_3)$ is disconnected. For example, consider the graph G' which is a circular ladder with six rungs. So G' is

two disjoint copies of C_6 with respective vertices labeled consecutively as $v_1, \dots, v_6$, and $w_1, \dots, w_6$, and additional edges $v_i \sim w_i$ for each i . Note that G' is bipartite, does not admit any folds, and every edge is contained in at least two copies of H_1 . Define $\eta(v_i) = \{i \bmod 3\}$ and $\eta(w_i) = \{(i+1) \bmod 3\}$ for all i . Then let τ be the restriction of η to the induced cycle on $\{v_1, \dots, v_6\}$. So $r(\tau) = 0$, and as before η and $\eta_{[l,j]}$ are in distinct connected components of $\text{Hom}(G', K_3)$ for any fixed pair $\{l, j\} \subset \{1, 2, 3\}$.

In the contrarian case of Q_3 , again every edge is contained in multiple copies of H_1 . The real issue, however, is that every 6-cycle in Q_3 is contained in either an induced copy of H_1 or an induced copy of $Q_3 \setminus \{v\}$ for some $v \in Q_3$. Both of these graphs fold to an edge, so $\text{Hom}(H_1, K_3)$ and $\text{Hom}(Q_3 \setminus \{v\}, K_3)$ are each connected. But then any distinct 0-cells $\tau_1, \tau_2 \in \text{Hom}(C_6, K_3)$ which extend to all of Q_3 must first extend to either H_1 or $Q_3 \setminus \{v\}$, where their extensions are necessarily in a single connected component.

5. $\text{Hom}(G(n, p), K_m)$

The topological connectivity of random hom-complexes has been studied previously under the guise of the Neighborhood Complex of $G(n, p)$, the Erdős–Rényi model for a random graph. $G(n, p)$ is the probability space of graphs on n vertices where each edge is inserted independently with probability $p = p(n)$. Kahle [9] showed that the connectivity of $\mathcal{N}(G(n, p))$ is concentrated between $1/2$ and $2/3$ of the expected value of the largest clique in $G(n, p)$, and also obtained asymptotic bounds on the number of dimensions with non-trivial homology. As we discussed earlier, $\mathcal{N}(G(n, p))$ is homotopy equivalent to $\text{Hom}(K_2, G(n, p))$. Here we take the opposite perspective and consider the random polyhedral complex $\text{Hom}(G(n, p), K_m)$.

The major benefit of utilizing $D(G)$ in a generalization of the Čukić–Kozlov Theorem is that k -cores have been well-studied in random graphs models. We say that $G(n, p)$ has property $\mathcal{P}$ with high probability if $\lim_{n \rightarrow \infty} \Pr[G(n, p) \in \mathcal{P}] = 1$. A property $\mathcal{P}$ is said to have a sharp threshold $\hat{p} = \hat{p}(n)$ if for all $\epsilon > 0$

$$\lim_{n \rightarrow \infty} \Pr[G(n, p) \in \mathcal{P}] = \begin{cases} 0 & \text{if } p \leq (1 - \epsilon)\hat{p} \\ 1 & \text{if } p \geq (1 + \epsilon)\hat{p} \end{cases}$$

When $p = c/n$ for constant $c > 0$, Pittel, Spencer and Wormwald [13] showed that the existence of a k -core in $G(n, p)$ has a sharp threshold $c = c_k$ for $k \geq 3$, and that asymptotically $c_k = k + \sqrt{k \log k} + O(\log k)$. Approximate values for small k are known, such as $c_3 \approx 3.35$, $c_4 \approx 5.14$, $c_5 \approx 6.81$. When $k = 2$, the existence of cycles has a one-sided sharp threshold at $c = 1$. For $c > 1$, indeed $\Pr[D(G(n, c/n)) \geq 2] \rightarrow 1$. However, for all $0 < c < 1$ there is a constant $0 < f(c) < 1$ for which $\Pr[D(G(n, c/n)) \geq 2] \rightarrow f(c)$. To simplify this issue, define c_k for all $k \geq 2$ by

$$c_k := \sup \left\{ c > 0 : \lim_{n \rightarrow \infty} \Pr[D(G(n, c/n)) \geq k] = 0 \right\}$$

In particular $c_2 = 0$, and for $k \geq 3$ these are precisely the sharp thresholds mentioned above.

By applying Theorem 2.8 to these thresholds, we then immediately obtain lower bounds on the topological connectivity of $\text{Hom}(G(n, p), K_m)$. For notational convenience, define $M(n, c, m) := \text{conn}[\text{Hom}(G(n, c/n), K_m)]$.

Theorem 5.1. *If $k \geq 2$ and $p = c/n$ with $c < c_{k+1}$, then for all $m \geq 3$*

$$\lim_{n \rightarrow \infty} \Pr[M(n, c, m) \geq m - k - 2] = 1$$

Theorem 2.8 does not require that the input graph be connected, and in fact $G(n, p)$ will be disconnected when $p = c/n$. Let the disjoint connected components of $G(n, p)$ be $G_0, G_1, \dots, G_t$, ordered from most vertices to least. For $c > 1$, with high probability G_0 is a giant component containing more than half the vertices, and G_i is either an isolated vertex, a tree, or a unicyclic graph for $i \geq 1$. So

$$\text{Hom}(G(n, c/n), K_m) = \text{Hom}\left(\bigsqcup_{i=0}^t G_i, K_m\right) = \prod_{i=0}^t \text{Hom}(G_i, K_m)$$

And thus

$$M(n, c, m) = \min_{0 \leq i \leq t} \{\text{conn}[\text{Hom}(G_i, K_m)]\}$$

For $G_i = \{v_i\}$, $\text{Hom}(G_i, K_m) = \Delta^2$, which is contractible. If a finite connected graph G_i is a tree, then it folds to a single edge and $\text{Hom}(G_i, K_m) \simeq \text{Hom}(K_2, K_m) \simeq S^{m-2}$. Finally, if G_i is a finite unicyclic graph, then it folds to its cycle C_n . If $n = 4$, then C_4 folds further to K_2 and $\text{Hom}(G_i, K_m) \simeq S^{m-2}$. Thus, with high probability we have

$$\text{Hom}(G(n, c/n), K_m) = (\Delta^2)^{t_1} \times (S^{m-2})^{t_2} \times \text{Hom}(G_0, K_m) \times \prod_{j=1}^{t_3} \text{Hom}(C_{n_j}, K_m)$$

where t_1 = the number of isolated vertices, t_2 = the number of trees and unicyclic components whose cycle is C_4 , $t_3 = t - (t_1 + t_2)$, and $n_j \neq 4$ for all j .

When $n_j = 3$, $\text{Hom}(C_{n_j}, K_m) = \text{Hom}(K_3, K_m)$, which is a wedge of $(m-3)$ -spheres [2]. Kozlov also computed the homology of $\text{Hom}(C_n, K_m)$ for all $n \geq 5$, $m \geq 4$ in [10]. In particular, for all $m \geq 4$:

$$\begin{aligned} \text{conn}[\text{Hom}(C_{2r+1}, K_m)] &= m-4 \quad \text{for } r \geq 1 \\ \text{conn}[\text{Hom}(C_{2r}, K_m)] &= m-3 \quad \text{for } r \geq 2 \end{aligned}$$

Note that the cases $n = 3$ and $n = 4$, which yield spheres, are consistent with these values. And the threshold for all small components to be isolated vertices is $p = \frac{\log n}{4n}$, so for $c < c_{k+1}$, $k \geq 2$, with high probability there is some $i \geq 1$ such that

$$\text{conn}[\text{Hom}(G_i, K_m)] \in \{m-3, m-4\}$$

This provides an upper bound for $M(n, c, m)$, and so when $m \geq 4$ we improve Theorem 5.1 to the following:

Theorem 5.2. *If $k \geq 2$ and $p = c/n$ with $c < c_{k+1}$, then for all $m \geq 4$*

$$\lim_{n \rightarrow \infty} \Pr[m-k-2 \leq M(n, c, m) \leq m-3] = 1$$

For fixed m , the lower bound decreases as c gets bigger, and the gap between upper and lower bounds becomes worse. We expect that $\text{conn}[\text{Hom}(G_0, K_m)]$ should decrease as $D(G_0)$ increases and the giant component becomes more highly connected. However, the lower bound in Theorem 2.8 is not tight in general. For example, the complete bipartite graph $K_{i,j}$ folds to an edge, so $\text{Hom}(K_{i,j}, K_m) \simeq S^{m-2}$. But $D(K_{i,j}) = \min\{i, j\}$, so Theorem 2.8 yields $\text{conn}[\text{Hom}(K_{i,j}, K_m)] \geq m - \min\{i, j\} - 2$. If $i, j \geq 2$, then this bound is not tight, and for large i, j it can be arbitrarily bad.

To sharpen these results, we turn to examining the chromatic number of $G(n, c/n)$. Similar to the thresholds for the appearance of k -cores, Achlioptas and Friedgut [1] showed that there is a sharp threshold sequence $d_k(n)$ such that for any $\epsilon > 0$,

$$c < (1 - \epsilon)d_k(n) \implies \lim_{n \rightarrow \infty} \Pr[\chi(G(n, c/n)) \leq k] = 1$$

$$c > (1 + \epsilon)d_k(n) \implies \lim_{n \rightarrow \infty} \Pr[\chi(G(n, c/n)) \geq k + 1] = 1$$

The convergence of the $d_k(n)$ remains an open problem, but Coja-Oghlan and Vilenchik [4] improved the lower bound on $\liminf_{n \rightarrow \infty} d_k(n)$ to within a small constant of the upper bound for $\limsup_{n \rightarrow \infty} d_k(n)$. In particular, for $o_k(1)$ a term which goes to 0 as k becomes large,

$$d_k^- := \liminf_{n \rightarrow \infty} d_k(n) \geq 2k \log k - \log k - 2 \log 2 - o_k(1)$$

$$d_k^+ := \limsup_{n \rightarrow \infty} d_k(n) \leq 2k \log k - \log k - 1 + o_k(1)$$

Then as k increases, the difference $|d_k^+ - d_k^-|$ approaches $2 \log 2 - 1 \approx 0.39$. And in the context of hom-complexes it is immediate that with high probability

$$\text{Hom}(G(n, c/n), K_m) = \emptyset \text{ for } c > d_m^+ \text{ and } \text{Hom}(G(n, c/n), K_m) \neq \emptyset \text{ for } c < d_m^-$$

For $m = 3$, we can then evaluate the connectivity of $\text{Hom}(G(n, c/n), K_3)$ for all $c > 0$, excluding the gap between d_3^- and d_3^+ .

Theorem 5.3. *For $1 \leq c < d_3^-$, $\text{Hom}(G(n, c/n), K_3)$ is disconnected with high probability.*

Proof of Theorem 5.3. By the definition of d_3^- and the existence of odd cycles with high probability for $c \geq 1$, $\chi(G(n, c/n)) = 3$. Then by Theorem 4.6 $\text{Hom}(G(n, c/n), K_3)$ is disconnected with high probability. $\square$

When $0 < c < 1$, all connected components are isolated vertices, trees, or unicyclic graphs with high probability. If there is a component which contains edges and does not fold to a single edge, then $\text{Hom}(G(n, c/n))$ will be disconnected. But if every connected component of $G(n, c/n)$ is an isolated vertex or folds to an edge, then $\text{Hom}(G(n, c/n), K_3)$ is connected. In the latter case, specifically

$$\text{Hom}(G(n, c/n), K_3) = (\Delta^2)^{t_1} \times \mathbb{T}^{t_2}$$

where t_1 = number of isolated vertices, and t_2 = the number of connected components which contain at least one edge. And $t_2 > 0$ with high probability for $p = c/n$, so the complex will not be contractible, and $M(n, c, 3) = 0$.

Theorem 5.4. *For $0 < c < 1$, $c' = \frac{1}{2} \log(1 - c) + \frac{c}{2} + \frac{c^2}{4} + \frac{c^4}{8}$,*

$$\lim_{n \rightarrow \infty} \Pr[M(n, c, 3) = -1] = 1 - e^{c'} \text{ and } \lim_{n \rightarrow \infty} \Pr[M(n, c, 3) = 0] = e^{c'}$$

Proof of Theorem 5.4. If G is a connected unicyclic graph whose cycle is C_l for $l \neq 4$, then G folds to C_l and $\text{Hom}(G, K_3) \simeq \text{Hom}(C_l, K_3)$, which is disconnected. So if $G(n, c/n)$ does not contain any cycle C_l for $l \neq 4$, then all components of $G(n, c/n)$ will be isolated vertices or will fold to an edge. For a fixed l , the number of l -cycles in $G(n, c/n)$ approaches a limiting Poisson distribution with mean $\frac{c^l}{2l}$, and so

$$\Pr[C_l \not\subset G(n, c/n) \text{ for all } l \neq 4] \rightarrow \exp\left\{-\frac{c^3}{6} - \sum_{l=5}^{\infty} \frac{c^l}{2l}\right\}$$

For $c < 1$, the sum in the exponent converges to $c' = \frac{1}{2} \log(1 - c) + \frac{c}{2} + \frac{c^2}{4} + \frac{c^4}{8}$. Thus $\Pr[M(c, n, 3) = 0] \rightarrow e^{c'}$ and $\Pr[M(c, n, 3) = -1] \rightarrow 1 - e^{c'}$. $\square$

6. FURTHER QUESTIONS

The biggest question left unanswered is that of bounding $\text{conn}[\text{Hom}(G_0, K_m)]$ from above. We expect that $\text{conn}[\text{Hom}(G_0, K_m)]$ remains close to $m - k - 2$ when this bound makes sense, that $M(n, c, m) = \text{conn}[\text{Hom}(G_0, K_m)]$, and that $M(n, c, m)$ is a non-increasing function for fixed m , with high probability. However, exhibiting non-trivial homology classes in $\text{Hom}(G, H)$ is difficult in general. In Section 4 we showed that disconnected components can be lifted via subgraphs, which can be viewed as lifting non-trivial 0-cycles, but there is no known analogous result for lifting higher dimensional non-trivial cycles.

We also expect that $M(n, c, m)$ increases monotonically when c is fixed and m is increasing, but this is not true in general for a fixed input graph. For instance, consider again our favorite counter example, Q_3 . We noted in Section 4 that $\text{conn}[\text{Hom}(Q_3, K_3)] \geq 0$. And by Theorem 2.8, $\text{conn}[\text{Hom}(Q_3, K_5)] \geq 0$. But for $m = 4$, there are colorings of Q_3 in which each color class is a pair of antipodal corners. In such a coloring, every vertex is adjacent to one vertex from each of the other three color class, so it represents an isolated 0-cell in $\text{Hom}(Q_3, K_4)$. Thus $\text{conn}[\text{Hom}(Q_3, K_4)] = -1$. In this example with Q_3 , the problem seems to occur when $m \leq D(G)$, and it is plausible that $\text{conn}[\text{Hom}(G, K_m)]$ is either increasing or non-decreasing for $m \geq D(G) + 1$, if G is a fixed graph which cannot be reduced via folds. This, however, remains an open question.

In the random setting, considering $m \leq D(G)$ presents a serious roadblock to evaluating $M(n, c, m)$ for all pairs of c and m . Since the $d_k^\pm$ grow much faster than the c_k , there are intervals where $D(G(n, c/n))$ is much larger than $\chi(G(n, c/n))$. For example, Molloy [12] pointed out that $c_{40} \approx 52.23$ while $d_{19} > 53.88$. So if $c = 53$, with high probability $D(G(n, c/n)) = 40$, but m can be as small as 19 before $\text{Hom}(G(n, c/n), K_m)$ becomes empty. When $m \leq D(G)$ the lower bound from Theorem 2.8 provides no information, and new methods are required.

A different direction altogether would be to indulge in a closer examination of precise numerical estimates on the Betti numbers and Euler characteristic of $\text{Hom}(G(n, c/n), K_m)$. Čukić and Kozlov's [7] work on cycles in hom-complexes makes the case that $m = 3$ a tantalizingly tractable place to start this type of investigation.

REFERENCES

- [1] Dimitris Achlioptas and Ehud Friedgut. A sharp threshold for k -colorability. *Random Structures Algorithms*, 14(1):63–70, 1999.
- [2] Eric Babson and Dmitry N. Kozlov. Complexes of graph homomorphisms. *Israel J. Math.*, 152:285–312, 2006.
- [3] Luis Cereceda, Jan van den Heuvel, and Matthew Johnson. Mixing 3-colourings in bipartite graphs. *European J. Combin.*, 30(7):1593–1606, 2009.
- [4] Amin Coja-Oghlan and Dan Vilenchik. Chasing the k -colorability threshold. In *2013 IEEE 54th Annual Symposium on Foundations of Computer Science—FOCS 2013*, pages 380–389. IEEE Computer Soc., Los Alamitos, CA, 2013.
- [5] Peter Csorba. *Non-Tidy Spaces and Graph Colorings*. PhD thesis, ETH Zürich, 2005.
- [6] Sonja Lj. Čukić and Dmitry N. Kozlov. Higher connectivity of graph coloring complexes. *Int. Math. Res. Not.*, (25):1543–1562, 2005.
- [7] Sonja Lj. Čukić and Dmitry N. Kozlov. The homotopy type of complexes of graph homomorphisms between cycles. *Discrete Comput. Geom.*, 36(2):313–329, 2006.
- [8] Alexander Engström. A short proof of a conjecture on the connectivity of graph coloring complexes. *Proc. Amer. Math. Soc.*, 134(12):3703–3705 (electronic), 2006.
- [9] Matthew Kahle. The neighborhood complex of a random graph. *J. Combin. Theory Ser. A*, 114(2):380–387, 2007.

- [10] Dmitry N. Kozlov. Cohomology of colorings of cycles. *Amer. J. Math.*, 130(3):829–857, 2008.
- [11] L. Lovász. Kneser’s conjecture, chromatic number, and homotopy. *J. Combin. Theory Ser. A*, 25(3):319–324, 1978.
- [12] Michael Molloy. A gap between the appearances of a k -core and a $(k + 1)$ -chromatic graph. *Random Structures Algorithms*, 8(2):159–160, 1996.
- [13] Boris Pittel, Joel Spencer, and Nicholas Wormald. Sudden emergence of a giant k -core in a random graph. *J. Combin. Theory Ser. B*, 67(1):111–151, 1996.
- [14] Carsten Schultz. Small models of graph colouring manifolds and the Stiefel manifolds $\text{Hom}(C_5, K_n)$. *J. Combin. Theory Ser. A*, 115(1):84–104, 2008.

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY